

Uemura plot as a certificate of two-dimensional character of superconducting transition for quisi-two-dimensional HTS

G.Sergeeva

National Science Center "Kharkov Institute of Physics and Technology",
Academicheskaya st.1, 61108, Kharkov, Ukraine
E-mail: gsergeeva@kipt.kharkov.ua

Abstract

For quisi-two-dimensional HTS in superconducting state the dimensional crossover, $3D \rightarrow 2D$ is studied. With using general properties of superconducting state for 2D systems the universal temperature dependence of the relation of the penetration lengths of magnetic field along axis $\hat{c}$, $\lambda^2(0)/\lambda^2(T/T_c)$, is found out. This dependence evidences about two-dimensional character of superconducting transition for quisi-two-dimensional HTS and leads to the Uemura plot for quisi-two-dimensional HTS.

1. Introduction. Quisi-two-dimensional (quisi 2D) or highly anisotropic underdoped HTS are exhibiting such out of the ordinary properties as "semiconducting-like" c-axis resistivity, $\rho_c(T)$, near T_c , big interval of two-dimensional superconducting fluctuations, $\Delta_{2D}^N \sim T_c$, and pseudo-gap states at $T \leq T^*$, where T^* is the temperature of charge ordering. These properties lead to the discussion of the association of superconducting transition with Berezinskii-Kosterlitz-Thouless (BKT) transition in CuO_2 planes at $T_{BKT} < T_{c0}$, where T_{c0} is the temperature of two-dimensional superconducting transition in the mean-field theory (see, for example, review [1] and the references there). Despite the observations of two-dimensional character of superconducting fluctuations and the evidences for a BKT transition have been reported in most of quisi 2D HTS [2,3], up to now the question whether a BKT transition is observable in bulk cuprates with taken into account interlayer coupling in these materials [4,5] is under the discussion.

Interlayer coupling can lead to the dimensional crossovers both as at $T \geq T_c$ ($2D \rightarrow 3D$), so in superconducting state near T_c ($3D \rightarrow 2D$) [4-6]. Theoretical model of such superconducting transition is well known [7-9]: at enough small probability t_c of charge tunnelling between CuO_2 planes transition has two-dimensional character with finite region Δ_{3D} of three-dimensional superconducting fluctuations. Anisotropy of exchange interactions in CuO_2 planes and along axis $\hat{c}$ in spin-fluctuational pairing model assumes big values of Δ_{2D}^N and T_{c0} and the distinction of temperatures $T_c \ll T_{c0}$. For quisi 2D HTS in ref. [10,11] it was shown that at $T > T_c$ two-dimensional character of superconducting fluctuations leads to the temperature dependence of the probability of charge tunnelling $t_c(T)$, and to the exasperation of semiconducting-like of $\rho_c(T)$ near T_c , so that $2D \rightarrow 3D$ crossover occurs at $T_c^0 > T_c > T_{BKT}$ before BKT transition. This point out to the two-dimensional character of superconducting transition, which develops under Kats scenario [7] at enough small value of probability $t_c(T_c^0)$:

$$T_c^0/\varepsilon_F < t_c(T_c^0), \quad (1)$$

where ε_F is Fermi energy, values T_c^0 and the temperature of three-dimensional transition in the mean field theory are the same order values. The interval Δ_{3D}^N of three-dimensional fluctuations in normal state can be found out from the measurements of resistivity $\rho_c(T)$

$$\Delta_{3D}^N \simeq T_c^0 - T_c \ll \Delta_{2D}^N \quad (2)$$

It is known that at $T < T_c$ back crossover $3D \rightarrow 2D$ occurs at T_{cr} which value depends on correlation length $\xi_c(T)$ along axis $\hat{c}$ [6, 12]. This paper is devoted to the studying of the dimensionality of the superconducting state of quisi 2D HTS and to the determination of the values T_{cr} . With using general properties of superconducting state for 2D systems the universal temperature dependence of the relation of the penetration lengths of magnetic field along axis $\hat{c}$, $\lambda^2(0)/\lambda^2(T/T_c)$, is found out.

2. Universal dependence of $T_c(\lambda^{-2}(0))$. The penetration lengths of magnetic field along axis $\hat{c}$, $\lambda(T)$ determines by London formula $\lambda(T) \simeq n_{s,3}^{-1/2}$, where $n_{s,3}$ is the three-dimensional superfluid density. The density $n_{s,3}$ evidently interconnects with two-dimensional superfluid density $n_{s,3} = n_s(T)\nu/l$, where ν is number of layers, l is the lattice constant. V.Pokrovskii shown that for quisi 2D HTS the penetration length $\lambda(T)$ and two-dimensional superfluid density $n_s(T)$ is connected by the expression

$$\lambda^2(0)/\lambda^2(T) = n_{s,3}(T)/n_{s,3}(0) = n_s(T)/n_s(0), \quad (3)$$

Here it will shown that for quisi 2D HTS (3) leads to universal dependence of $T_c(\lambda^{-2}(0))$,

which was called "Uemura plot" and was discovered at the measurements of muon relaxation rate [14].

For plane system (3) can be written as ratio $\rho_s(T/T_c)/\rho_s(0) = n_s(T/T_c)/n_s(0)$, where $\rho_s(T/T_c)$ is dimensionless hardness and satisfy to universal dependence [15]

$$\rho_s(T/T_c) = \exp\left(-\frac{Te^{-1}}{T_c\rho_s(T/T_c)}\right) \quad (4)$$

The decision of (4) was received in ref.[15]: $\rho_s(0) = 1$, and at $T = T_c$

$$\rho_s(T/T_c)|_{T=T_c} = e^{-1}, \quad (5)$$

Expressions (3)-(4) results to universal character of temperature dependence of ratio

$$\lambda^2(0)/\lambda^2(T/T_c) = \rho_s(T/T_c)/\rho_s(0) = \exp\left(-\frac{Te^{-1}\lambda^2(T/T_c)}{T_c\lambda^2(0)}\right), \quad (6)$$

and simple relation between values $\lambda^2(T)$ and $n_s(T)$ at $T = T_c$ and at $T = 0$:

$$\lambda^2(0)/\lambda^2(T_c) = n_s(T)/n_s(0) = e^{-1} \quad (7)$$

Using this relation and Kosterlitz-Thouless-Nelson formula [16]

$$k_B T_c = \frac{h^2}{32\pi m} n_s(T_c), \quad (8)$$

we can receive the universal relation between the density $n_s(0)$ at $T = 0$ and transition temperature T_c :

$$T_c = \frac{h^2 e^{-1}}{32k_B \pi m} n_s(0), \quad (9)$$

where k_B is Boltzman constant. For two-dimension superconductor the role of effective penetration length acts magnetic screening length

$$L_s(T) = \frac{mc^2}{2\pi n_s(T)e^2}, \quad (10)$$

which is connected with the bulk London penetration length

$$L_s = 2d^{-1}\lambda^2, \quad (11)$$

where d - thickness of CuO_2 plane.

It is seen from (7-11) that the transition temperature is proportional to $\lambda^{-2}(0)$

$$T_c = k\lambda^{-2}(0), \quad (12)$$

where

$$k = \frac{c^2 h^2 e^{-1} d}{64k_B \pi^2 e^2}, \quad (13)$$

depends only from universal constants and from thickness of CuO_2 plane. The temperature of dimensional crossover, $T_{cr} < T_c$, depends on correlation length $\xi_c(T)$, and can be determined at the measurement of penetration length as the boundary of two-dimensional region, where at $T > T_{cr}$ the measurement values of $\lambda^2(0)/\lambda^2(T/T_c)$ are diverging from universal relation (6). Knowing T_{cr} let us to determine the interval $\Delta_{3D}^S \approx (T_c - T_{cr})$ of three-dimensional fluctuations at $T < T_c$. Full interval is equal

$$\Delta_{3D} = \Delta_{3D}^N + \Delta_{3D}^S = T_c^0 - T_{cr} \quad (14)$$

does not depend on the exactness of the measurement of T_c , and determines as the difference between the temperatures of two dimensional crossovers: in normal state, T_c^0 , and in superconducting state, T_{cr} . So, for $La_{1.85}Sr_{0.15}CuO_4$ the measurements of resistivity $\rho_c(T)$ [17] and of the penetration length [18] let us to determine the temperature $T_c^0 \sim 41.5K$, and $T_{cr} \sim 26K$. This lead to the value $\Delta_{3D} \sim 15.5K$. We see that the region Δ_{3D} of three-dimensional superconducting fluctuations is finite in the concrete, that evidences about the two-dimensional character of superconducting transition.

Thus, here it is shown that expression (12), which was found out at the measurements of muon relaxation rate [14] for quisi 2D HTS, really is universal and is the sequence of general consistent pattern (4), (8) of superconducting state for two-dimensional systems and must to fulfil for HTS with two-dimensional superconducting state at $T < T_{cr}$. This means that for quisi 2D HTS superconducting transition has two-dimensional character and develops on Kats scenario [7], and at $T < T_{cr}$ the temperature dependence of $\lambda^2(0)/\lambda^2(T/T_c)$ also must to be universal (6).

The author would like to thank Dr.Kabanov V. for the reference on the V.L.Pokrovskii paper [13].

References

1. S.L.Cooper, K.E.Gray, in "Physical properties of high temperature superconductors" ed.: Donald M.Ginsberg,IY p.61-188 (1994).
2. S.Martin, A.T.Fiory, R.M.Fleming, G.P.Espinoza, and A.S.Cooper, Phys.Rev.Lett. **62**, 677 (1989).
- 3.D.H.Kim, A.M.Goldman, J.H.Kang, R.T.Kampwirth, Phys.Rev. **B40**, 8834 (1989).
4. Y.Matsuda, S.Komiyama, T.Onogi, T.Terashima, K.Shimura, and Y.Bando , Phys.Rev.**B48**, 10498 (1993).
5. Y.Matsuda, S.Komiyama, T.Terashima, K.Shimura, and Y.Bando, Phys.Rev.Lett. **69**, 3228 (1992).
6. Z.Tesanovic, L.Xing, L.Bulaevskii, O.Li, and Suenaga, Phys.Rev.Lett.**69**, 3563 (1992).
7. E.I.Kats, JEF (Russian),**56**, 1675 (1969).

8. Vik.Dotsenko and M.V.Feigelman, JETP,**83**, 345 (1982).
9. L.Bulaevskii, Usp.Fiz.Nauk (Russian), **116**, 449 (1975).
10. G.Sergeeva, Physica C,**341-348**, 181 (2000).
11. G.G.Sergeeva, Low Temp. Phys.,**26**, 453 (2000); cond-mat/0009212.
12. T.Schneider and H.Keler, Phys.Rev.**B47**, 5915 (1993).
13. V.L.Pokrovskii, Pis'ma JETP, (Russian) **47**, 539 (1988).
14. Y.J.Uemura, V.J.Emery, A.R.Moodenbaugh et al. Phys.Rev.**B38**, 909 (1988); Phys.Rev.Lett.**64**, 2082 (1990).
15. Patashinskii A.Z., Pokrovskii V.L. Fluctuational theory of phase transition (Russian, Moscow, Nauka) 1982.
16. B.I.Halperin and David R.Nelson, J. Low Temp.Phys. **36**, 599 (1979).
17. T.Ito, H.Takagi, S.Ishibashi et al. Nature, **350**, 596 (1991).
18. G.Aeppli and R.J.Cava, E.J.Ansaldo et al. Phys.Rev.**B35**, 7129 (1987).